

Solution of the Anderson impurity model via the functional renormalization group

Simon Streib, Aldo Isidori, and Peter Kopietz
*Institut für Theoretische Physik, Universität Frankfurt,
 Max-von-Laue Strasse 1, 60438 Frankfurt, Germany*
 (Dated: November 7, 2012)

We show that the functional renormalization group is a numerically cheap method to obtain the low-energy behavior of the Anderson impurity model describing a localized interacting electron coupled to a bath of conduction electrons. Our approach uses an external magnetic field as flow parameter, partial bosonization of the transverse spin fluctuations, and frequency-independent interaction vertices which are fixed by Ward identities. We calculate the quasi-particle residue and the spin susceptibility in the particle-hole symmetric case and obtain excellent agreement with the Bethe ansatz results for arbitrary strengths of the interaction.

PACS numbers: 72.15.Qm, 71.27.+q, 71.10.Pm

The Anderson impurity model (AIM) describes a localized interacting electron in contact with a bath of non-interacting conduction electrons¹. The model was first introduced in the context of material science for describing the emergence of local moments in metals and has been studied for half a century by means of a variety of methods². In the past decade, renewed attention has been drawn to the AIM because of its experimental realization in quantum dot systems in contact with a metallic environment. Moreover, the solution of the AIM is one of the fundamental steps in the so-called dynamical mean-field theory³. In practice, this step is often carried out using Wilson's numerical renormalization group (NRG), which yields numerically controlled results for the thermodynamics and the spectral properties⁴. In the 1980s the thermodynamics of the AIM has also been obtained exactly via the Bethe ansatz⁵. For later reference, we quote here the Bethe ansatz results for the spin and charge susceptibilities in the particle-hole symmetric case⁶,

$$\pi\Delta\chi_s = \sqrt{\frac{2}{\pi u}} e^{\pi^2 u/8} \int_0^\infty dx e^{-x^2/(2u)} \frac{\cos(\pi x/2)}{1-x^2}, \quad (1a)$$

$$\pi\Delta\chi_c = \sqrt{\frac{2}{\pi u}} e^{-\pi^2 u/8} \int_0^\infty dx e^{-x^2/(2u)} \frac{\cosh(\pi x/2)}{1+x^2}. \quad (1b)$$

Here $u = U/(\pi\Delta)$, where U is the interaction between two localized electrons with opposite spin, and Δ is the hybridization energy to the conduction band, in the limit where the latter has infinite bandwidth and constant density of states. For $u \gtrsim 2$ the charge susceptibility χ_c becomes exponentially small, while the spin susceptibility $\chi_s = (4T_K)^{-1}$ is proportional to the inverse of the Kondo temperature $T_K = \Delta(\pi u/2)^{1/2} e^{-\pi^2 u/8 + 1/(2u)}$.

Although Eqs. (1a, 1b) can be confirmed numerically using the NRG (up to small deviations due to the finite width of the conduction band in the NRG calculations), it would be useful to have an alternative and numerically less expensive method to obtain the correct low-energy physics of the AIM at strong coupling, which is dominated by the exponentially small Kondo temperature. In this work we show that this can be achieved by means of the functional renormalization group (FRG) method⁷. In

the past few years, many authors have studied the AIM using different versions and truncations of the FRG⁸⁻¹⁴, but none of these works succeeded in reproducing the correct strong coupling behavior of the AIM. We show below that this problem can be solved by means of a rather simple truncation of the FRG hierarchy involving only frequency-independent vertices, provided that the transverse spin fluctuations are properly bosonized and the interaction vertices are fixed by Ward identities.

For simplicity, we focus on the particle-hole symmetric AIM. After integrating out the conduction electrons, we arrive at the following Euclidean action for the Grassmann field $d_\sigma(\tau)$ describing the localized electron,

$$S = - \int_\omega \sum_\sigma G_{0,\sigma}^{-1}(i\omega) \bar{d}_{\omega\sigma} d_{\omega\sigma} + U \int_0^{1/T} d\tau n_\uparrow n_\downarrow. \quad (2)$$

Here T is the temperature (we set $k_B = \hbar = 1$ and focus on the zero temperature limit below), $\int_\omega = T \sum_\omega$ denotes summation over fermionic Matsubara frequencies $i\omega$, $n_\sigma = \bar{d}_\sigma(\tau) d_\sigma(\tau)$ represents the number of localized electrons with spin $\sigma = \uparrow, \downarrow$, and $d_{\omega\sigma} = \int_0^{1/T} d\tau e^{i\omega\tau} d_\sigma(\tau)$. Assuming that the conduction band has infinite width and constant density of states, the inverse Green function is $G_{0,\sigma}^{-1}(i\omega) = i\omega - \xi_\sigma + i\Delta \text{sgn} \omega$, where in the particle-hole symmetric case the energy of the localized electron relative to the chemical potential is $\xi_\sigma = -U/2 - \sigma H$, and H is the external magnetic field in units of energy.

Our strategy is inspired by the field-theoretical renormalization group approach recently developed by Edwards and Hewson¹⁵, who noticed that for large magnetic fields the AIM can be studied perturbatively, even for $U \gg \Delta$, as long as U/H is sufficiently small. Following Ref. [15], let us therefore use the external magnetic field H as a flow parameter for the FRG. Unfortunately, using this magnetic field cutoff scheme we have not been able to reproduce the strong-coupling physics of the AIM if we formulate the FRG exclusively in terms of fermionic degrees of freedom. The problem is that the resulting fermionic vertices exhibit a rather strong frequency dependence which cannot be neglected. To avoid this problem, we partially bosonize the action (2)

by decoupling the interaction in terms of a complex boson field χ representing transverse spin fluctuations. It turns out to be advantageous, however, to bosonize only a part of the interaction term, while retaining a fermionic parametrization for the rest. Therefore, we write the interaction in Eq. (2) as $Un_\uparrow n_\downarrow = U_\parallel n_\uparrow n_\downarrow - U_\perp s_+ s_-$, where $U = U_\parallel + U_\perp$ and the Grassmann bi-linears $s_+ = \bar{d}_\uparrow d_\downarrow$ and $s_- = \bar{d}_\downarrow d_\uparrow$ represent the usual spin ladder operators. Only the transverse part $-U_\perp s_+ s_-$ is now bosonized by introducing a complex boson field χ via a usual Hubbard-Stratonovich transformation^{10,11}. We choose $U_\perp$ such that it vanishes for $H \rightarrow \infty$, and that it reduces to the bare interaction U for $H \rightarrow 0$. Therefore we set $U_\perp^{-1} = U^{-1} + R_H$ and require that $R_0 = 0$ and $R_\infty = \infty$. Note that R_H plays the role of a regulator function for the bosonic sector of the theory which switches on the interaction between transverse spin fluctuations. The proper choice of R_H will be discussed in more detail below.

It is now straightforward to write down formally exact FRG flow equations for the irreducible vertices of our mixed boson-fermion model^{7,10}. For our purpose, it is sufficient to neglect the frequency dependence of all one-line irreducible vertices with more than two external legs. We also neglect the contribution from charge- and longitudinal spin fluctuations to the FRG flow¹⁶. The resulting FRG flow equation for the self-energy $\Sigma_\sigma(i\omega)$ of the localized electron is shown graphically in Fig. 1; the corresponding analytic expression reads

$$\begin{aligned} \frac{\partial \Sigma_\sigma(i\omega)}{\partial H} &= \int_{\omega'} \sum_{\sigma'} \dot{G}_{\sigma'}(i\omega') \Gamma_4^{\bar{d}_\sigma \bar{d}_{\sigma'} d_{\sigma'} d_\sigma} \\ &+ \int_{\bar{\omega}} \dot{F}_\perp(i\bar{\omega}) \Gamma_4^{\bar{d}_\sigma d_\sigma \bar{\chi} \chi} + \int_{\omega'} \left[\dot{G}_{-\sigma}(i\omega') F_\perp(i\sigma(\omega - \omega')) \right. \\ &\quad \left. + G_{-\sigma}(i\omega') \dot{F}_\perp(i\sigma(\omega - \omega')) \right] \Gamma_3^{\bar{d}_\uparrow d_\downarrow \chi} \Gamma_3^{\bar{d}_\downarrow d_\uparrow \bar{\chi}}. \end{aligned} \quad (3)$$

The single-scale propagators in our cutoff scheme are

$$\dot{G}_\sigma(i\omega) = -\sigma G_\sigma^2(i\omega), \quad \dot{F}_\perp(i\bar{\omega}) = -F_\perp^2(i\bar{\omega}) \partial R_H / \partial H, \quad (4)$$

where $G_\sigma(i\omega)$ is the exact fermionic Green function and $F_\perp(i\bar{\omega})$ is the exact propagator of the spin fluctuation field χ . These propagators can be expressed in terms of the corresponding irreducible self-energies $\Sigma_\sigma(i\omega)$ and $\Pi_\perp(i\bar{\omega})$ via the Dyson equations $G_\sigma^{-1}(i\omega) = G_{0,\sigma}^{-1}(i\omega) - \Sigma_\sigma(i\omega)$ and $F_\perp^{-1}(i\bar{\omega}) = U_\perp^{-1} - \Pi_\perp(i\bar{\omega})$. We denote bosonic Matsubara frequencies by $i\bar{\omega}$. The right-hand side of the flow equation (3) depends on the irreducible four-point vertices $\Gamma_4^{\bar{d}_\sigma \bar{d}_{\sigma'} d_{\sigma'} d_\sigma}$ and $\Gamma_4^{\bar{d}_\sigma d_\sigma \bar{\chi} \chi}$ at vanishing external frequencies, and on the three-point vertices $\Gamma_3^{\bar{d}_\uparrow d_\downarrow \chi} = \Gamma_3^{\bar{d}_\downarrow d_\uparrow \bar{\chi}} \equiv \gamma$. The superscripts indicate the types of external legs attached to the vertices.

Our goal is to derive flow equations for the spin-dependent part M of the fermionic self-energy at vanishing frequency and the quasi-particle residue Z , which are defined via the low-energy expansion

$$\Sigma_\sigma(i\omega) = U/2 - \sigma M + (1 - Z^{-1})i\omega + \mathcal{O}(\omega^2). \quad (5)$$




FIG. 1. (Color online) Graphical representation of the FRG flow equation (3) for $\Sigma_\uparrow(i\omega)$ (shaded circle). Solid and wavy lines represent full fermion and boson propagators, respectively. For each type of line, the slash insertions denote the corresponding single-scale propagators in Eq. (4). The dot on the left-hand side represents a derivative with respect to H .

In order to obtain M and Z from the solution of the flow equation (3) we need to know the bosonic self-energy $\Pi_\perp(i\bar{\omega})$ and three different vertices: $\Gamma_4^{\bar{d}_\sigma \bar{d}_{\sigma'} d_{\sigma'} d_\sigma}$, $\Gamma_4^{\bar{d}_\sigma d_\sigma \bar{\chi} \chi}$, and γ . One could write down additional flow equations for these vertices, but these depend on higher order vertices and it is not clear how to truncate this infinite hierarchy. We shall therefore adopt a different strategy which uses Ward identities and skeleton equations to obtain a closed system of flow equations for M and Z .

First of all, we note that the three-point vertex at vanishing frequencies can be related to the self-energy parameter M via the Ward identity

$$\Gamma_3^{\bar{d}_\uparrow d_\downarrow \chi} = \Gamma_3^{\bar{d}_\downarrow d_\uparrow \bar{\chi}} \equiv \gamma = b/(h + u_\perp \arctan b), \quad (6)$$

which can be derived for our bosonized theory following the work by Koyama and Tachiki^{17,18}. Here $u_\perp = U_\perp/(\pi\Delta)$ and $b = h + m$, with $m = M/\Delta$ and $h = H/\Delta$.

Next, let us derive an alternative flow equation for M which does not explicitly involve the four-point vertices. To this end we consider the longitudinal spin-susceptibility $\chi_\parallel = \partial s / \partial H$, where $s = \langle n_\uparrow - n_\downarrow \rangle / 2$ is the local magnetic moment (which we normalize such that $s = 1/2$ corresponds to a fully magnetized state). The local moment is then connected to the self-energy parameter M via the Friedel sum rule²,

$$\pi s = \arctan(h + m) = \arctan b. \quad (7)$$

Taking the derivative with respect to h we obtain

$$\partial m / \partial h = \chi_\parallel / \rho - 1, \quad (8)$$

where $\rho = [\pi\Delta(1 + b^2)]^{-1}$ is the exact density of states at vanishing energy. To determine $\chi_\parallel$ we use the Ward identity^{2,19}

$$\chi_\parallel / \rho = 1/Z + \rho \Gamma_\perp, \quad (9)$$

where $\Gamma_\perp$ is the exact interaction vertex between two electrons with opposite spin at vanishing frequencies. In our partially bosonized theory we have

$$\Gamma_\perp = \gamma^2 U_\perp / [1 - U_\perp \Pi_\perp(0)] + \Gamma_4^{\bar{d}_\uparrow \bar{d}_\downarrow d_\downarrow d_\uparrow}. \quad (10)$$

But the three-point vertex and the fermionic four-point vertex turn out to be mutually related via a skeleton equation. With the help of standard functional methods⁷ it is indeed straightforward to show that, at finite frequencies,

$$\Gamma_3^{\bar{d}_\uparrow d_\downarrow \chi}(\omega + \bar{\omega}, \omega; \bar{\omega}) = 1 - \int_{\omega'} G_\downarrow(i\omega') G_\uparrow(i\omega' + i\bar{\omega}) \times \Gamma_4^{\bar{d}_\uparrow \bar{d}_\downarrow d_\downarrow d_\uparrow}(\omega + \bar{\omega}, \omega'; \omega, \omega' + \bar{\omega}). \quad (11)$$

Using the above relation to express $\Gamma_4^{\bar{d}_\uparrow \bar{d}_\downarrow d_\downarrow d_\uparrow}$ in terms of $\Gamma_3^{\bar{d}_\uparrow d_\downarrow \chi} \equiv \gamma$ at vanishing external frequencies, we find for the longitudinal susceptibility²⁰,

$$\frac{\chi_\parallel}{\rho} = \frac{1}{Z} + \frac{1}{1+b^2} \frac{b}{\arctan b} \left[\frac{b}{h} - \frac{1}{Z} \right], \quad (12)$$

where we have used the Ward identity (6) to eliminate the vertex γ . Substituting the expression (12) into Eq. (8), we obtain a flow equation for m which depends only on the self-energy parameters m and Z .

Finally, to close our system of flow equations we approximate the bosonic self-energy $\Pi_\perp(i\bar{\omega})$, appearing implicitly in Eq. (3), by

$$\pi\Delta\Pi_\perp(i\bar{\omega}) = \gamma P_\perp(|\bar{\omega}|/(Z\Delta), b \operatorname{sgn} \bar{\omega}), \quad (13)$$

where the dimensionless function

$$P_\perp(x, b) = \frac{\ln[1 + x/(1 - ib)]}{(x - 2ib)(1 + x/2 - ib)} - \frac{2i \arctan b}{x - 2ib} \quad (14)$$

is obtained from the skeleton equation¹⁰ for $\Pi_\perp(i\bar{\omega})$. The prefactor in Eq. (13) is determined by demanding that $\Pi_\perp(0)$ satisfy the Ward identity^{15,17}

$$\Pi_\perp(0)/[1 - U_\perp\Pi_\perp(0)] \equiv \chi_\perp = s/H \quad (15)$$

relating the transverse spin-susceptibility $\chi_\perp$ to the local moment s . Using the Friedel sum rule (7) it is easy to see that our expression (12) for the longitudinal susceptibility is consistent with Eq. (15) for $\chi_\perp$, in the sense that for $H \rightarrow 0$ both susceptibilities approach the same limit χ_s . We point out that Eqs. (6) and (15) imply that $a \equiv [1 - U_\perp\Pi_\perp(0)] = h/(h + u_\perp \arctan b) > 0$, which guarantees that there is no magnetic instability.

We have now obtained a closed system of flow equations for $m = M/\Delta$ and Z . For simplicity, we expand the function $P_\perp(x, b)$ in Eq. (13) to linear order in x , which is consistent with the low-energy expansion (5) in the fermionic sector. After straightforward algebra we arrive at the flow equation $h\partial Z/\partial h = \eta Z$ for the quasi-particle residue Z , with

$$\eta = hZ\gamma^2 u_\perp \operatorname{Im} \int_0^\infty \frac{dx}{(1 - ib + x)^2} \frac{c}{(a + cx)^2} + hZ\gamma^2 u_\perp^2 \frac{\partial r}{\partial h} \operatorname{Re} \int_0^\infty \frac{dx}{(1 - ib + x)} \frac{2c}{(a + cx)^3}. \quad (16)$$

Here $r = \pi\Delta R_H$ is the dimensionless regulator function of the boson propagator, and

$$c = (u_\perp \gamma)(2ib)^{-1} [(1 - ib)^{-2} - \arctan b/b]. \quad (17)$$




FIG. 2. (Color online) Logarithmic plot of the bosonic regulator $r = \pi\Delta R_H$ as a function of the magnetic field $h = H/\Delta$, for $U/(\pi\Delta) = 1$ (solid line) and $U/(\pi\Delta) = 8$ (dashed line). The regulator is linear in h both at small and large h , with a change in the prefactor from α ($h \lesssim 1$) to β ($h \gtrsim 1$). The inset shows the U -dependence of the coefficients α and β .

The above system of flow equations can be easily solved numerically once the bosonic regulator function $r = \pi\Delta R_H$ has been specified. The simplest choice of a linear magnetic field dependence of the bosonic regulator ($r \propto h$, in agreement with the magnetic field regulator in the fermionic sector) does not lead to satisfactory results. After testing many regulators, we found that the best results can be obtained with a regulator of the form

$$r = (\alpha - \beta)h/(1 + h^2) + \beta h, \quad (18)$$

which is linear in h both for small and large h , but exhibits a change in slope from α to β at $h \approx 1$, as shown in Fig. 2. Comparing our FRG results at weak coupling, $U \ll \Delta$, with the known perturbative expansion for Z and χ_s , we also realized that the coefficients α and β are weakly dependent functions of the dimensionless bare interaction $u = U/(\pi\Delta)$. Their functional dependence is well described by rational functions of low degree. Hence, we introduce the following ansatz,

$$\alpha = \alpha_0/(\alpha_2 + u^2), \quad \beta = \beta_1 u/(\beta_2 + u^2), \quad (19)$$

where the four parameters α_0 , α_2 , β_1 , and β_2 can be determined by matching the FRG results for Z and χ_s at weak coupling (e.g., for $u = 0.1$) with the corresponding values obtained in the perturbative expansion at $\mathcal{O}(u^2)$, and by imposing that the Wilson ratio R is equal to 2 for two values of u in the Kondo regime (e.g., we impose $R = 2$ for $u = 4$ and $u = 8$). Note that in the AIM the expression for the Wilson ratio reads

$$R \equiv \frac{\chi_s(U)/\chi_s(0)}{Z^{-1}(U)/Z^{-1}(0)} = \pi\Delta\chi_s Z = \frac{2\chi_s}{\chi_s + \chi_c}, \quad (20)$$

where the last equality follows from the Yamada-Yosida Ward identities^{19,21}. At large U , charge fluctuations are




FIG. 3. (Color online) Inverse quasi-particle residue Z^{-1} and spin susceptibility χ_s as functions of $U/(\pi\Delta)$ for $H = 0$. The FRG results for Z^{-1} (squares) and χ_s (circles) are compared with the corresponding Bethe ansatz results (solid and dashed lines, respectively). As a reference, we also show Z^{-1} calculated within second order perturbation theory (dotted line). The inset shows the FRG (diamonds) and Bethe ansatz (solid line) results for the Wilson ratio R .

strongly suppressed and the low-energy physics of the AIM is effectively the same as in the Kondo model. In this regime the charge susceptibility χ_c is therefore negligible and the Wilson ratio takes the Kondo-model value $R = 2$. For the parametrization of α and β in Eq. (19), we find $\alpha_0 \approx 0.36$, $\alpha_2 \approx 1.4$, $\beta_1 \approx 0.19$, and $\beta_2 \approx 5.4$. The U -dependence of α and β is shown in the inset of Fig. 2.

Having fixed the regulator in the bosonic sector, we can now solve our FRG flow equations numerically, with a minimal computational cost. Choosing the initial magnetic field H_0 sufficiently large, we may use as initial conditions for M and Z the Hartree-Fock values $M_0 = U/2$ and $Z_0 = 1$, which are exact in the limit $H_0 \rightarrow \infty$. The FRG results for Z and χ_s , in the limit of vanishing magnetic field, are shown in Fig. 3, where in the inset we also show the corresponding Wilson ratio. The remarkable agreement of our approach with the Bethe ansatz results for arbitrary large values of the interaction shows that the present FRG truncation, supplied with exact Ward identities, is able to capture the exponentially small Kondo scale in the low-energy quasi-particle resonance.

In Fig. 4 we show, for $u = U/(\pi\Delta) = 5$, the magnetic field dependence of Z , s , and γ . In the magnetization curve $s(H)$ one can clearly identify the two energy scales characterizing the strong coupling regime of the AIM, namely the bare Coulomb interaction U and the width $Z\Delta$ of the Kondo resonance. Indeed, reducing the external magnetic field H from a large initial value, at $H \approx U$ we start observing a slow decrease in the magnetic moment from the fully magnetized value $s = 1/2$. The decrease in s becomes then faster when $H \approx Z\Delta$, until the




FIG. 4. (Color online) Inverse quasi-particle residue Z^{-1} (dotted-dashed line, scale on the right), local magnetic moment s (solid line), and three-point vertex γ (dashed line) as functions of the external magnetic field H , for $U/(\pi\Delta) = 5$.

magnetization eventually vanishes for $H \ll Z\Delta$. The presence of two different energy scales is also reflected in the non-monotonic behavior of the three-point vertex $\gamma(H)$. Starting from the bare value $\gamma = 1$ at large H , the vertex becomes initially larger than unity for $H \approx U$ and later decreases again, approaching (in the Kondo regime) the value $1/u < 1$ for $H \lesssim Z\Delta$.

In summary, we have developed an FRG approach to the AIM which uses the external magnetic field H as a physical flow parameter, and where the fermionic interaction U is partially decoupled via a bosonic field describing the transverse spin fluctuations. We have truncated the exact hierarchy of FRG flow equations keeping only the frequency-independent parts of the interaction vertices and expanding the fermionic and bosonic self-energies to linear order in frequency. With an extensive use of Ward identities, we have been able to express all the relevant vertices in terms of self-energy parameters, thereby avoiding further approximations in the flow equations for the vertices. Comparing our results with the exact Bethe ansatz solution at vanishing magnetic field, we have shown that the present FRG approach is able to reproduce the exponential U -dependence of the width of the Kondo resonance. Moreover, the use of the magnetic field H as a physical cutoff for the FRG flow equation gives access to the non-trivial H -dependence of physical observables such as the local magnetic moment. Possible extensions of the present method include the study of the non-symmetric AIM, the use of finite frequency terms in the fermionic self-energy in order to describe the high-energy features of the spectral function, and a generalization of our approach to non-equilibrium situations relevant for the transport properties of quantum dots.

We thank A. C. Hewson for useful discussions and for

making his unpublished notes on the AIM available to us. This work was supported by the DFG via FOR 723.

-
- ¹ P. W. Anderson, Phys. Rev. **124**, 41 (1961).
² A. C. Hewson, *The Kondo Problem to Heavy Fermions*, (Cambridge University Press, Cambridge, 1993).
³ A. Georges, G. Kotliar, W. Krauth, and M. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).
⁴ K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975); R. Bulla, T. Costi, and T. Pruschke, *ibid.* **80**, 395 (2008).
⁵ A. M. Tsvelick and P. B. Wiegmann, Adv. Phys. **32**, 453 (1983).
⁶ V. Zlatić and B. Horvatić, Phys. Rev. B **28**, 6904 (1983).
⁷ P. Kopietz, L. Bartosch, and F. Schütz, *Introduction to the Functional Renormalization Group*, (Springer, Berlin, 2010).
⁸ R. Hedden, V. Meden, T. Pruschke, and K. Schönhammer, J. Phys.: Cond. Mat. **16**, 5279 (2004).
⁹ C. Karrasch, R. Hedden, R. Peters, T. Pruschke, K. Schönhammer, and V. Meden, J. Phys.: Cond. Mat. **20**, 345205 (2008).
¹⁰ L. Bartosch, H. Freire, J. J. R. Cardenas, and P. Kopietz, J. Phys.: Cond. Mat. **21**, 305602 (2009).
¹¹ A. Isidori, D. Roosen, L. Bartosch, W. Hofstetter, and P. Kopietz, Phys. Rev. B **81**, 235120 (2010).
¹² S. G. Jakobs, M. Pletyukhov, and H. Schoeller, Phys. Rev. B **81**, 195109 (2010).
¹³ H. Freire and E. Corrêa, J. Low Temp. Phys. **166**, 192 (2012).
¹⁴ M. Kinza, J. Ortloff, J. Bauer, and C. Honerkamp, arXiv:1210.0763.
¹⁵ K. Edwards and A. C. Hewson, J. Phys.: Condens. Matter **23**, 045601 (2011).
¹⁶ Note that in our magnetic field cutoff scheme the part $U_{\parallel}n_{\uparrow}n_{\downarrow}$ of the interaction which is not bosonized depends on the cutoff H , which therefore does not appear exclusively in the Gaussian part of the propagator. We can avoid this technical complication by writing $U_{\parallel}n_{\uparrow}n_{\downarrow} = U_{\parallel}[(n_{\uparrow} + n_{\downarrow})^2 - (n_{\uparrow} - n_{\downarrow})^2]/4$ and decoupling the quadratic terms by means of additional Hubbard-Stratonovich fields representing charge- and longitudinal spin fluctuations.
¹⁷ T. Koyama and M. Tachiki, Prog. Theor. Phys. Supplement **80**, 108 (1984).
¹⁸ S. Streib, A. Isidori, and P. Kopietz, unpublished.
¹⁹ P. Kopietz, L. Bartosch, L. Costa, A. Isidori, and A. Ferraz, J. Phys. A: Math. Theor. **43**, 385004 (2010).
²⁰ Since we are neglecting the frequency dependence of all vertices, we need to introduce a correction factor in Eq. (11). This can be fixed by demanding that for $\hbar \rightarrow 0$ the resulting longitudinal susceptibility $\chi_{\parallel}$ in Eq. (12) approach the same limit as the transverse susceptibility $\chi_{\perp}$ in Eq. (15).
²¹ K. Yamada, Prog. Theor. Phys. **53**, 970 (1975); K. Yosida and K. Yamada, *ibid.* **53**, 1286 (1975); K. Yamada, *ibid.* **54**, 316 (1975).